

TRAUMA SURVIVORS INNER ROLE ATOM: A CLINICAL MAP FOR POST-TRAUMATIC GROWTH**Scott Giacomucci, MSS, LCSW, CTTS, CET III, CP, PAT, TSI-TL****Abstract**

The treatment of Post-Traumatic Stress Disorder (PTSD) and other trauma-related maladies requires psychotherapists to be equipped with a dependable clinical map that can act as a compass through the difficulties of trauma therapy. The Therapeutic Spiral Model (TSM), a clinically sophisticated and research-supported adaptation of classical psychodrama, comes equipped with a comprehensive clinical map called the Trauma Survivor's Inner Role Atom (TSIRA) which emphasizes safety, containment, and strengths. TSM has been utilized in over 30 countries around the world. The TSIRA provides a guide to intrapsychic structural change conceptualized in the simplicity of role theory while drawing from continued developments neuroscience research. The TSIRA offers a triune map beginning with *prescriptive roles* to build strengths, connection, accurate observation, containment, and safety. The *trauma roles* offered by TSM's intrapsychic trauma triangle are explored only after the prescriptive roles have been established, with the clinical functions of each demonstrated. And finally, the *transformative roles*, the internal manifestation of post-traumatic growth, emerge and are integrated as a completion of the clinical map's three spirals. The implementation of this inner role atom as a clinical map prevents retraumatization while providing emotional regulation to protagonists and the group keeping them within their window of tolerance.

Keywords: *Therapeutic Spiral Model, Trauma, Post-Traumatic Stress Disorder, Post-Traumatic Growth, Neuroscience, clinical map*

INTRODUCTION

The Trauma Survivors' Inner Role Atom (TSIRA) emerged spontaneously from the development of the Therapeutic Spiral Model (TSM), a research-supported (Gow & McVea, 2006; Greensberg, 2013; Hudgins, Drucker, & Metcalf, 2000; Hudgins & Toscani, 2013; Perry, Saby, Wenos, Hudgins, & Baller, 2016) and clinically modified form of classical psychodrama that emphasizes safety, containment, and strengths in the treatment of Post-Traumatic Stress Disorder (PTSD) (Hudgins, 2017b; Hudgins & Toscani, 2013). The TSIRA is the very core of TSM and influences everything that a TSM practitioner does, including team meetings before and after workshops, introductions, warm-ups, sociometry, psychodrama, and sharing. It is meant to be an internal role atom, or template, that provides a structure for healthy personality functioning using non-pathologizing role theory and the latest in neurobiology research. It facilitates the safety needed to establish a therapeutic alliance and group cohesion while keeping clients in their window of tolerance and transforming the internalized trauma-based roles into post-traumatic growth.

Working with trauma can be a difficult and intimidating process, for both the client and therapist. The possibility of retraumatizing, or reinforcing the effects of trauma, is real (Cozolino 2010; Hudgins & Toscani, 2013; Schore, 2012; van der Kolk, 2014). The very nature of trauma lends itself towards re-experiencing and reenacting the past in the present moment. It is essential that we allow our interventions to be guided by a clinical map, rather than only working "intuitively" with our traumatized clients or "following the [traumatized] protagonist". The lasting effects of trauma – insecure attachments, hyperarousal, persistent negative thoughts and mood states, avoidance, and re-experiencing symptoms (American Psychiatric

Trauma Survivor's Inner Role Atom

Association, 2013) create fertile ground for the dance of transference and counter-transference between client and therapist. At times it will be difficult to discern between intuition and counter-transference. For these reasons, it is in the best interest of both client and therapist that we employ a clinical map to safely direct our work.

The simplest way to describe the TSIRA is using a visual of a spiral with three strands – prescriptive roles, trauma-based roles, and transformative roles. The first strand represents prescriptive roles which focus on developing the ability for non-judgmental observation, containment, and strengths. The term *prescriptive* is used to reflect that these roles are directives from a professional and are necessary for the change to occur, just like a prescription from a medical doctor. The second spiral symbolizes the internalization of the trauma. And the transformation that emerges between the interaction of prescriptive and trauma-based roles is represented by the final strand of the spiral. The TSIRA provides a template with intervention steps that target the development of specific psychological functions necessary for healthy functioning after trauma (Hudgins, 2017a).

PRESCRIPTIVE ROLES: AN EXPANSION OF THE SELF

J.L. Moreno offers us a simple conceptualization of the self as being the sum of all of the roles that we play (1946). The TSIRA is a diagram that offers a template for a healthy self. Each of the three spiral strands are expanded into the 18 labeled roles of the TSIRA, which is an internal role atom, or diagram of internal roles necessary for a healthy personality. TSM works primarily with internal roles rather than interpersonal roles, which means that each TSM drama takes place within the brain or the self of an individual. Instead of following the protagonist, like

Trauma Survivor's Inner Role Atom

in classical psychodrama, the prescriptive roles of the TSIRA are first enrolled within any TSM drama to ensure safety. The TSIRA *prescribes* roles relating to restoration, containment, and observation before moving into any traumatic material. These 8 roles prevent retraumatization and keep protagonists within their window of tolerance – the threshold where healing can take place (Siegel, 2011). The prescriptive roles are the operational definition of spontaneity (Hudgins, 2017b) and allow protagonists to respond in a new way to their trauma (Hudgins & Toscani, 2013).

Function	Prescriptive Roles
A. Observation	1. Observing Ego 2. Client Role
B. Containment	3. Containing Double 4. Body Double 5. Manager of Defenses
C. Restoration/Strength	6. Intrapsychic Strengths 7. Interpersonal Strengths 8. Transpersonal Strengths

Table 1. Prescriptive roles and functions (Toscani & Hudgins, 1995)

Prescriptive Roles of Observation

The first function, observation, is concretized by TSM's role of the *Observing Ego*, a term originally used by Freud (1932), which is the part of self that can accurately observe the self without blame, shame, or judgement. This is a cognitive role associated with the left hemisphere and cognitive functions of the brain that helps to keep the protagonist present and provide accurate labeling for emotionally charged events which are likely stored in nonverbal memory (Lawrence, 2015). In TSM workshops, this role is often concretized with cards of some kind (animal, angel, Buddha, art work, etc.) which are placed around the room. The Observing Ego, with similarities to classical psychodrama's mirror role (Moreno, Blomkvist, & Rutzel,

Trauma Survivor's Inner Role Atom

2000), allows the protagonist to see herself within the context of the scene - helping to piece together fragmented parts of traumatic memory into a meaningful narrative.

The second TSM role of observation is the *Client Role*, which is seldom used in clinical practice, except when working with Dissociative Identity Disorders (Hudgins & Toscani, 2013). The Client Role is a role like the Observing Ego which observes the psychodrama, and later helps the protagonist integrate her experience before leaving the session. This role provides the client with a method of concretizing the part of self with more personal work to do – having the goal of providing both observation and containment for the client in-between sessions.

Prescriptive Roles of Containment

The containment functions are operationalized by TSM's unique roles of the *Containing Double*, *Body Double*, and the *Manager of Defenses*. The Containing Double is a role that stays with the protagonist at all times, offering statements anchoring them in the present moment and expanding or containing feeling or thinking, when clinically appropriate. The Containing Double adapts based on the needs of each protagonist. For a protagonist with overwhelming feelings, the Containing Double would contain the feelings while helping to label internal experience; but for a protagonist prone to intellectualizing or overthinking, the Containing Double would contain the thinking while helping him access his feelings and physical sensations. One might say that it serves as the corpus callosum, connecting the left and right hemispheres of the brain and providing a balance between cognition and emotion (Hug, 2013). Multiple research studies on the Containing Double have been produced which support its efficacy (Hudgins & Drucker, 1998; Hudgins, Drucker, & Metcalf, 2000).

Trauma Survivor's Inner Role Atom

On the other hand, the Body Double holds the role of a healthy body for the protagonist and offers statements around grounding, breathing, eye-contact, and posture (Burden & Ciotola, 2001; Carnabucci & Ciotola, 2013). This role of the Body Double becomes especially helpful in reducing dissociation for clients who do not feel safe in their bodies and clients with eating disorders. The Body Double reconnects the trauma survivor with awareness of her own body thus strengthening vertical integration and providing grounding (Lawrence, 2011). This process stimulates the use of both proprioception and interoception for the protagonist, which are core therapeutic elements of other trauma therapies (van der Kolk, 2014) - including Somatic Experiencing (Levine, 2008, 2010; Payne, Levine, Crane-Godreau, 2015), Sensorimotor Psychotherapy (Ogden, Minton, & Pain, 2006), and Yoga-Based Practices (Schmalzl, Powers, & Blom, 2015). Often in clinical practice, the Body Double and Containing Double become integrated into one dynamic role.

The Manager of Defenses is a role that is clinically necessary when a protagonist begins to demonstrate psychological defenses that might prevent them from achieving the clinical contract in the psychodrama. Defenses are seen as attempts to maintain safety and are honored for their function at the time of the trauma. The protagonist is invited to consider if the defense is still needed today. Each of these roles of containment are demonstrated in the following clinical vignette from an inpatient treatment group at Mirmont Treatment Center.

In contracting for a psychodrama, a client states her goal of learning how to love herself while demonstrating signs of dissociation. The director invites her to choose someone to play the role of her Containing Double and Body Double (which are sometimes combined into one role). She walks about the room with the double beside her making statements such as "I can

Trauma Survivor's Inner Role Atom

take a deep breath and stay present," "when I notice myself dissociating, I can feel my feet on the floor and find the safety here in the room," and "I am deepening my love for myself right now by doing this work". At which point the director asks the protagonist to concretize her defense of dissociation using an object in the room by asking "who can hold your dissociation? You do not appear to need it in this moment". The protagonist then selects a group member to play the role of the Manager of Defenses and consciously gives the concretized dissociation to the Manager of Defenses. This provides a visualization of the dissociation as being externalized and contained, allowing the protagonist to remain more fully present. Role-reversing the protagonist into the role of the Manager of Defenses offers her an embodied and empowering experience of managing her defense. The utilization of the defense then becomes a choice instead of an unconscious decision or a habitual pattern. This is a clear example of spontaneity in action as a protagonist with a chronic pattern of dissociation, through the prescriptive roles of containment, was able to have "a new response to an old situation" (Moreno, 1953, pg. 336).

Prescriptive Roles of Restoration/Strength

The final psychological function of the prescriptive roles is restoration and strengths. There are three types of strengths; *intrapsychic*, *interpersonal*, and *transpersonal*. Intrapsychic strengths are those within the individual, such as courage, determination, and hope. Interpersonal strengths may be supportive people or strengths that involve more than just the self. Examples include a good friend, a teacher, or the ability to trust others. The last type of strength is transpersonal, which simply means that it is beyond human. This could mean God, a higher power, elements from myths or legends, music, nature, a healthy ancestral legacy, or a greater purpose in life.

Trauma Survivor's Inner Role Atom

In TSM clinical practice, these strengths are first concretized (usually with scarves) in a safety structure called the “Circle of Strengths” which involves group members identifying their own, and each other's strengths, in a ritual of acknowledging self and others. The Circle of Strengths, strengthens group connections, acknowledges strengths, and creates a physical circle – which serves as both a container for the trauma and a stage for the psychodrama. Furthermore, this action structure helps the group gradually enter the space of surplus reality, which Moreno (1965) defines as a mode of subjective experience beyond reality which is enriched through imagination (1965, pg. 212-213).

Integrating Neuroscience and TSM's Prescriptive Roles

When a protagonist volunteers, offers her trauma-related topic to the group, and states her goal for the work, she is tapping into both the memory of the trauma and the associated neural network. TSM's prescriptive roles, which are enrolled into the drama before any trauma-based or antagonist roles are incorporated, help to renegotiate the emotional context of the traumatic memory by activating different affective systems and providing a felt sense of safety after the memory has been stimulated. These strength-based roles, are most likely to initiate the PLAY, CARE, and SEEKING social-emotional systems, as described by Panksepp & Biven (2012), which in effect provide a renegotiation and recontextualization of the traumatic memory (Levine, 2015).

In role-reversals with nurturing prescriptive roles, a TSM practitioner utilizes the role-reciprocity between the protagonist and the prescriptive role to activate both the protagonist's spontaneity and her CARE system as they role-train compassion and care for self. Panksepp &

Trauma Survivor's Inner Role Atom

Biven (2012) write that experiencing feelings of PANIC/GRIEF in others (even through a role-player!) is one of the most powerful triggers of the CARE system – thus highlighting one of the neurobiological mechanisms of role-reciprocity in psychodrama. Furthermore, activating the protagonist's CARE system (in the role-reversal) will inhibit her GRIEF system (pg. 285), effectively strengthening the renegotiation of the memory's emotional context.

The fact that another human plays the role of these roles is neurobiologically significant. The interaction with a role (and thus the group member playing the role) activates the protagonist's ventral vagal nerve ("Smart Vagus") and social engagement system providing emotional regulation (Porges, 2017). On the other hand, the auxiliary roles' interactions with the protagonist throughout the drama provide an experience of consistent attunement which is also associated with emotional regulation and feelings of security, safety, self-esteem, confidence, and connection (Fishbane, 2007). Furthermore, because "the 'self' is largely a construction of the prefrontal cortex or the thinking mind" (Dayton, 2015, pg. 111), which is often frozen when trauma becomes activated, the experience of role reversing into any other role allows the protagonist to concretize and physically see herself - helping to keep the prefrontal cortex of the brain actively stimulated.

Neurobiologically, TSM's prescriptive roles help prevent uncontrolled regression and stimulate intrapsychic change by providing clinically necessary interventions that foster bilateral hemispheric (left-right brain) integration, vertical integration (Siegel 2011), mind/body integration, and social integration within the group. Interestingly, addiction researcher Bruce Alexander (2011) emphasizes that the primary cause of addictions is a lack of psychosocial integration, often resulting from trauma (Maté, 2008).

Trauma Survivor's Inner Role Atom

Daniel Siegel (2011), who coined the term “interpersonal neurobiology”, defines integration as the key element upholding mental health and defines eight different types of integration. He states that “our task is to find the impediments to the eight domains of integration and liberate the mind’s natural drive to heal – to integrate mind, brain, and relationships” (2011, pg. 76). I was struck by the parallel between Siegel and Zerka Moreno’s words on the process of healing – “protagonists themselves do the healing. My task is to find and touch that autonomous healing center within, to assist and direct the protagonist to do the same” (2012, pg. 504). The Therapeutic Spiral Model, with the guidance of its clinical map, effectively walks the protagonist through removing the barriers to integration and the mind’s innate ability to heal itself – what Zerka labels the autonomous healing center.

“Integration is the goal, not catharsis” (Hug, 2013, pg. 129). Or, to express it in classical psychodrama terms, a catharsis of integration must follow a catharsis of abreaction (Hollander, 1969; Moreno, 1953). It is only fitting that the *integration* of all eight of TSM’s prescriptive roles culminates into one of TSM’s most important transformative roles, the Appropriate Authority, which provides a protagonist with the empowerment to stop repeating the internalization of the trauma triangle.

THE TRAUMA TRIANGLE: THE INTERNALIZATION OF TRAUMA

The TSIRA offers a simple, and non-pathologizing conceptualization of the internalization of trauma through TSM’s unique trauma triangle. The trauma triangle was an evolution of Karpman’s (1968) interpersonal drama triangle of victim, perpetrator, and rescuer. In one’s experience of trauma however, there was no rescuer, otherwise the trauma would not

Trauma Survivor's Inner Role Atom

have occurred. So TSM teaches that a trauma survivor unconsciously internalizes the roles of *Victim, Perpetrator, and Abandoning authority* (Hudgins & Toscani 2013; Toscani & Hudgins, 1995). These three trauma-based roles are the TSM operational definition of PTSD symptomology in action.

While an individual is actively on this intrapsychic trauma triangle, they are significantly inhibited from connecting with others. The primary benefit to getting off the trauma triangle is for the purposes of engaging in more meaningful and productive relationships with self and others. Similarly, when one is engaged in a response to danger (fight, flight, or freeze), their social engagement and attachment systems are strongly suppressed making connection with others nearly impossible.

Triangulated Roles: Victim, Perpetrator, & Abandoning Authority

The Victim role holds the woundedness and the story of the trauma, but often feels too unsafe to speak it, resulting in hyperarousal and isolation from others. Simultaneously, the Perpetrator role holds the personal power of the survivor, but uses it to continuously perpetrate self-harm, often years after the trauma is over. There are many ways that the internalized perpetrator role manifests, mostly as intrusions, flashbacks, and symptoms of reexperiencing. The final role, unique to TSM, is the Abandoning Authority which presents through the abandonment of self in behaviors such as addictions, avoidance, dissociation, not standing up for or taking care of self. The final PTSD symptom cluster, Negative Cognitions and Moods, appears to exist in the middle of the triangle, being related to all three of the trauma

Trauma Survivor's Inner Role Atom

roles. The internalized trauma role-taking and the presentation of trauma symptomology mirror each other as parallel processes using different language to frame the same phenomenon.

These three internal roles, Victim, Perpetrator, and Abandoning Authority create a triangulation of role reciprocity. TSM theory conceptualizes the trauma as living within the survivor in terms of these roles which can be thought of as the living introjections of the spoken and unspoken messages from the perpetrator and abandoning authority at the time of the trauma. Although the actual trauma is over, it lives within the survivor and is reexperienced through the surplus reality of flashbacks, night terrors, negative cognitions and feeling states, avoidance, dissociation, and insecure attachments (American Psychiatric Association, 2013).

TSM Psychodrama uses the surplus reality of prescriptive roles to alter the way in which the trauma lives within the protagonist, which is another form of surplus reality. The interaction of the prescriptive roles with the trauma-based roles is exactly what creates the intrapsychic change according to TSM theory. TSM defines the prescriptive roles as the definition of spontaneity in action (Hudgins, 2017b) which, when interacting with the trauma-based roles, allows the protagonist to respond in a new, adequate way instead of resorting to the repetitive trauma triangle patterns. The alchemy of prescriptive roles interacting with trauma-based roles is what creates transformative roles; this will be depicted in a future vignette.

TRANSFORMATIVE ROLES: AN OPERATIONAL DEFINITION OF POST-TRAUMATIC GROWTH


Trauma Survivor's Inner Role Atom

Post-Traumatic Growth refers to phenomenon of positive transformation that is often experienced after a traumatic life event (Calhoun & Tedeschi, 2014). The TSIRA's transformative roles are the operational definition of post-traumatic growth in action and embodied in the simplicity of role theory. The TSIRA's transformative roles include eight labelled roles organized on the three poles of transformative functions - autonomy, integration, and correction. These functions can be conceptualized of as the opposite sides of the trauma triangle roles constituting role transformations from abandonment to integration, victimhood to autonomy, and perpetration to correction (Giacomucci, 2018).

Perpetrator

Roles of Correction

Good-Enough Parent
Good-Enough Partner
Good-Enough Friend



Victim

Roles of Autonomy

Sleeping-Awakening Child
Change Agent
Manager of Healthy
Defenses

Abandoning Authority

Roles of Integration

Appropriate Authority
Ultimate Authority

Figure 1. TSM Trauma Triangle Role Transformations (Giacomucci, 2018).

This diagram depicts the TSM transformative triangle (heart-shaped) as an evolution of the TSM trauma triangle with the alignment of trauma-based roles and the corresponding TSM Transformative roles and functions.

Transformative roles of Integration

Trauma Survivor's Inner Role Atom

One of the most important transformative roles on the TSIRA clinical map is the *Appropriate Authority*, which is necessary to help remove one's self from cycling around the internal trauma triangle (Hudgins & Toscani 2013). The Appropriate Authority is an internal role that intervenes in the repetition of continued abandonment, victimization, and perpetration of the self. This is an integrated part of self that halts negative introjections stemming from the experience of trauma as the trauma survivor begins to claim back her personal power in an appropriate manner.

TSM's other role of integration, the *Ultimate Authority*, is the integration of all 8 of the transformative roles having been internalized, enacted in the protagonist's intrapsychic world, then her interpersonal world, and finally out in the world. This role is, in a spiritual sense, awakening to the fact that one is a co-creator and co-responsible for mankind (Moreno, 2012).

Transformative Roles of Autonomy

The TSIRA transformative roles, enacting the function of autonomy, help the protagonist to free herself from the past and create a new future. Even someone with a relatively healthy personality uses psychological defense mechanisms while navigating and interacting with the complexity of the world. *The Manager of Healthy Defenses* offers one the power to *choose* to use a defense appropriately rather than be a victim to the chronic, repetitive use of defenses in response to a lack of safety that many trauma survivors experience.

The *Sleeping-Awakening Child* is another role unique to TSM. Many trauma survivors indicate that they feel as though they have lost their innocence, spontaneity, creativity, or inherent goodness. The Sleeping-Awakening Child role reframes these beliefs and offers a new

Trauma Survivor's Inner Role Atom

construct; this is the role that holds all of the innocence, goodness, uniqueness, creativity, and spontaneity. It was never lost or taken, it simply went to sleep at the time of the trauma and waits for the protagonist to make her life safe enough to be awoken (Hudgins 2017a). It is a truly beautiful moment in a TSM psychodrama to experience an auxiliary play the role of the Sleeping-Awakening Child as the protagonist awakens this part of self, and in doing so, taps into a source of inner goodness.

The *Change Agent* is a role that is always present and is most responsible for the change that takes place within the psychodrama. Once internalized, this role can be utilized by the protagonist to be an agent of change in her own life and the world. This role often manifests in different ways from drama to drama taking on various different faces.

Transformative Roles of Correction

The Transformative roles of corrective connection, which are *Good-Enough Parents*, *Good-Enough Significant Other*, and *Good-Enough Spirituality*, are significant in their ability to provide protagonists with corrective emotional experiences that have the power to repair the negative influence of prior experiences (Alexander, 1946; Cozolino, 2014). Often, the clinical contract for a TSM psychodrama is to have an experience of developmental repair from the original trauma. In psychodramas of developmental repair, the protagonist is guided through the stages of the TSIRA, starting with prescriptive roles, moving onto defenses and into the trauma-based roles, at which point the transformative roles emerge spontaneously. A brief example of this type of drama is depicted in the following section.

**TSM PSYCHODRAMA OF POST-TRAUMATIC GROWTH: RE-EXPERIENCING WITH
DEVELOPMENTAL REPAIR**

The following vignette is a depiction of an TSM psychodrama in an addiction and trauma inpatient group at Mirmont Treatment Center in Lima, PA. First, the group establishes safety and warm-ups to the psychodrama by participating in an exercise concretizing their *Observing Egos* using *TSM Animal Cards* and placing them visibly around the room. Following, group members take turns acknowledging each other's strengths, and in doing so, concretize a large circle of strengths which serves as a stage. John is sociometrically chosen by the group as protagonist voicing that he would like to heal from his experience of re-occurring physical abuse from his deceased father. He expresses that he always craved to hear his father say, "I love you." John communicates that for most of his life, he has been very critical of himself and has low self-worth which he believes is related to his experience of abuse. After the goal for the psychodrama is stated, the scene begins to emerge - always beginning with prescriptive roles. The director, based on his clinical assessments and experience of John, begins to *prescribe* necessary roles in addition to the roles emerging spontaneously.

Prescriptive Role Scene: Resourcing the Protagonist

The first scene of this drama includes John connecting to a *Containing/Body Double*, the *courage* and *determination* to face his trauma, his *resilience*, his *best friend* as an interpersonal support, and transpersonal strengths as the mythical *Hercules* and an *angel*. At one point, he becomes discouraged at the thought of facing his trauma. The director brings him to concretize his feeling of inadequacy with a scarf which John chooses to ask *Hercules* to hold (Hercules now

Trauma Survivor's Inner Role Atom

holds both the transpersonal strength and Manager of Defenses roles). He is brought to role-reverse with his Observing Ego card and sees himself surrounded by strengths and supportive roles which help both to increase his confidence and see his adult-self more accurately.

Hercules encourages him to embark on this hero's journey into his trauma!

Trauma Triangle Scene: Renegotiating the Trauma

The director enrolls John's wounded, *younger self* (Victim role) into the scene to provide him with an experience of being protected by his strengths, nurturing himself from the adult role, and being nurtured from the child role. John promises not to abandon himself again (role transition from Abandoning Authority to Appropriate Authority), but to stay present and continue doing the work necessary to heal. The next scene involves *John's father* (Perpetrator) entering the drama with a loud, demeaning, and critical attitude towards John. Because TSM is an intrapsychic model, the role of his father is labeled as an introjected part of self which lives within him long after the trauma and even after his father's death. John, supported by his strengths, shouts back at his father and stands up for himself. A contained, though physically-involved struggle takes place between John and his father until John pushes his father out of the scene. Of course, prior to the scene, the group contracts for the necessary safety to conduct the physically demanding scene that takes place.

A Scene of Transformations: Post-Traumatic Growth

While forcing the introjects from his father out of his brain, he firmly states his worth as a human being and tells his father that the abuse was wrong and unfair. Next, as we move into the transformative scene, two group members are enlisted to hold the roles of these two

Trauma Survivor's Inner Role Atom

themes that spontaneously emerged as a direct result of the interaction between prescriptive roles and trauma-based roles - *worthiness* (Sleeping-Awakening Child) and *fairness/justice* (Appropriate Authority). John connects again to the younger self and acknowledges his goodness and worthiness aloud. Without the messages from the internalized Perpetrator role, he finds it much easier to acknowledge his goodness. The role of worthiness (Sleeping-Awakening Child) becomes much more active and full of life!

Finally, the director, knowing John's beliefs about the afterlife, invites the angel role to bring John to interact with his father's spirit. The angel explains that, in the afterlife, an individual's defects and humanness are shed as they become a pure spirit again. John begins a dialogue with the spirit of his father who apologizes for his abusive actions and commends his son for such brave personal growth work. The spirit of John's father tells him that he loves him as John embraces his father and both begin to experience tears of joy.

Clinical Processing of the Psychodrama

Through this psychodrama, John had an embodied experience of a loving father that he may have never had in his life. Just as traumatic experiences change us, so do positive experiences - whether they are psychodramatic or not. For John, the psychodrama was a multisensory experience like any other. The presence of his abusive father earlier in the scene helped John access the associated traumatic neural network, while the prescriptive roles helped him move from his "stuckness" or immobilization into sympathetic activation, creating a new ending to the scene. John moved through the subsystems outlined in the Polyvagal theory in the order in which they evolutionarily developed – from the most primitive immobilization

Trauma Survivor's Inner Role Atom

(freeze) system, to the mammalian sympathetic system (fight/flight response), and finally to social engagement (Porges, 2017). This provides his nervous system with an experience of completing the survival responses unavailable to it at the time of the childhood abuse - resulting in a reframing of the internalized trauma, both in terms of intrapsychic role organization (Hudgins & Toscani 2013) and neurobiological wiring (Cozolino, 2010, 2014). In TSM clinical practice, the director never role-reverses the protagonist into a victim role (wounded self) until she has demonstrated her capacity to rescue the victim from the adult ego state first. This helps to prevent retraumatization and serves as an indicator that the protagonist will not get stuck in the victim role once role-reversed.

The corrective experience of being in relationship with a *Good-Enough Father* during the psychodrama helps facilitate a repair in his attachment schema, as well as his own introjections and self-talk, which appear to stem from his abusive father. This would not be possible if it weren't for the brain's innate mechanisms of neuroplasticity and continued learning. The brain undergoes synaptic and structural changes directly related to new social interactions and experiences (Cozolino, 2010, 2014; Siegal, 2012). Quite literally, experiences change our brain and have the power to reverse the impact of earlier traumatic events. Furthermore, the internalized "imprint of trauma doesn't sit in the verbal, understanding part of the brain" says Bessel van der Kolk, "but in much deeper regions – amygdala, hippocampus, hypothalamus, brain stem – which are only marginally affected by thinking and cognition" (as cited in Wylie, 2004, pp. 30-41). Thus underscoring one of the reasons why experiential therapies are the treatment of choice for individuals with PTSD and trauma-related issues (Dayton, 2015; Hudgins 2017a; van der Kolk, 2014).

CONCLUSION

The Therapeutic Spiral Model, through its use of psychodramatic enactments of developmental repair, provides participants with embodied corrective experiences directly related to the trauma or neglect that was experienced earlier in life. The application of TSM's clinical map, the TSIRA, acts as a holding environment for clients, containing them within their window of tolerance while renegotiating their trauma and facilitating a completion of the central nervous system's survival responses (Levine, 2008, 2010; Porges, 2017). Through the use of surplus reality, clients have the opportunity to have their developmental needs met, reverse the effects of trauma, and actively change their brain. Utilizing neuroplasticity, changes are materialized by accessing the original traumatic neural network with safety, and through role-playing change the experience, or create a new experience. TSM psychodrama allows participants to play the roles of transformation and post-traumatic growth in the safety of a psychodrama, effectively role-training them to hold the roles in other arenas of their lives.

TSM is an intrapsychic model and its clinical map is oriented on inner roles. TSM integrates object relations, attachment, and role theory to teach that interpersonal relationships, positive and negative, are internalized. The founders of the model came to realize that one needed to do their own inner work before they could take change out into their relationships and ultimately into the world. While TSM psychodramas are always staged within the brain of the protagonist and thus focus on inner roles, attention is given to interpersonal

Trauma Survivor's Inner Role Atom

roles and relationships in both the warm-up and sharing phases of the psychodrama and TSM's six safety structures.

Unfortunately, we cannot change the fact that our clients have experienced trauma. However, through the proper application of TSM's clinical map, we can help our clients transform their internalized trauma into post-traumatic growth. As our clients internalize their own intrapsychic Change Agent role, they become empowered to change not just their inner role atom (TSIRA), but also their social atom. Through their development of the Ultimate Authority role of integration, they become initiators in the world and claim co-responsibility for all of mankind (Moreno, J., 1947; Moreno, Z., 2012).

REFERENCES

- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders: DSM-5*. Washington, D.C: American Psychiatric Association.
- Burden, K. & Ciotola, L. (2001). *The Body Double: An Advanced Clinical Action Intervention Module in the Therapeutic Spiral Model to Treat Trauma*. Retrieved from <http://www.healing-bridges.com/psychodrama.html>.
- Calhoun, L.G. & Tedeschi, R.G. (2014). *The Handbook of Post-Traumatic Growth: Research and Practice*. New York, NY: Psychology Press.
- Carnabucci K. & Ciotola, L. (2013). *Healing Eating Disorders with Psychodrama and Other Action Methods: Beyond the Silence and the Fury*. London, UK: Jessica Kingsley Publishers.
- Cozolino, L. J. (2010). *The Neuroscience of Psychotherapy: Building and Rebuilding the Human Brain (2nd Ed)*. New York: W.W. Norton & Company.
- Cozolino, L. J. (2014). *The Neuroscience of Human Relationships: Attachment and the Developing Social Brain (2nd Ed)*. New York: W.W. Norton & Company.
- Dayton, T. (2015). *NeuroPsychodrama in the Treatment of Relational Trauma: A Strength-based, Experiential Model for Healing PTSD*. Deerfield Beach, FL: Health Communications, Inc.
- Fishbane, M.D. (2007). Wired to Connect: Neuroscience, Relationships, and Therapy. *Family Process* 46: 395-412.
- Freud, S. (1932). *The dissection of the psychical personality. Standard edition 22, 67-80*. London: Hogarth Press.

Trauma Survivor's Inner Role Atom

Giacomucci, S. (2018). Transforming the TSM Trauma Triangle: From Roles of Post-Traumatic Stress to Roles of Post-Traumatic Growth. *Therapeutic Spiral International Newsletter*. Charlottesville, VA. Retrieved from TherapeuticSpiralModel.com.

Greenberg, L. S. (2013). Anchoring the therapeutic spiral model into research on experiential psychotherapies. In K. Hudgins & F. Toscani (Eds.), *Healing world trauma with the therapeutic spiral model*. (pp. 132-148) London, UK: Jessica Kingsley Publishers.

Gow, K. & McVea, C. (2006) Healing a Mother's Emotional Pain: Protagonist and Director Recall of a Therapeutic Spiral Model (TSM) Session. *Journal of Group Psychotherapy, Psychodrama & Sociometry*, 59(1), 3-22.

Hollander, C. E. (1969). *A process for psychodrama training: The Hollander psychodrama curve*. Denver, CO: Snow Lion Press.

Hudgins, K. (2017a). A Simple Clinical Action Map to Heal PTSD: The Therapeutic Spiral Model's Intrapsychic Role Atom. *Applied Neuropsychotherapy*. Accessed July 22nd from: www.neuropsychotherapist.com/a-simple-clinical-action-map-to-heal-ptsd/

Hudgins, K. (2017b) PTSD Unites the World: Prevention, Intervention and Training in The Therapeutic Spiral Model. In C.E. Stout and G. Want (eds). *Why Global Health Matters: Guidebook for Innovation and Inspiration*. CreateSpace Independent Publishing Platform

Hudgins, M.K, & Drucker, K. (1998). The Containing Double as Part of the Therapeutic Spiral Model for Treating Trauma Survivors. *International Journal of Action Methods*, 51(2), 63-

Trauma Survivor's Inner Role Atom

Hudgins, M. K, Drucker, K & Metcalfe, K (2000). The Containing Double: A Clinically Effective Psychodrama Intervention for PTSD. *British Journal of Psychodrama and Sociodrama*, 15(1): 58-77

Hudgins, K. & Toscani, F. (2013). *Healing World Trauma with The Therapeutic Spiral Model: Stories From the Frontlines*. London, UK: Jessica Kingsley Publishers.

Hug, E. (2013). A Neuroscience perspective on trauma and action methods. In K. Hudgins, & F. Toscani (Eds.), *Healing world trauma with the therapeutic spiral model*. London, UK: Jessica Kingsley Publishers.

Karpman, S. (1968). Fairy Tales and Script Drama Analysis. *Transactional Analysis Bulletin* 7(26) 39-43

Lawrence, C. (2011). *The architecture of mindfulness: Integrating the Therapeutic Spiral Model and interpersonal neurobiology*. Retrieved from <http://www.therapeuticspiral.com>

Lawrence, C. (2015). The Caring Observer: Creating Self-Compassion Through Psychodrama. *The Journal of Psychodrama, Sociometry, and Group Psychotherapy* 63(1): 65-72

Levine, P.A. (2008). *Healing Trauma: A Pioneering Program for Restoring the Wisdom of Your Body*. Berkeley, CA: North Atlantic Books.

Levine, P.A. (2010). *In an Unspoken Voice: How the Body Releases Trauma and Restores Goodness*. Berkeley, CA. North Atlantic Books.

Levine, P.A. (2015). *Trauma and Memory: Brain and Body in a Search for the Living Past*. Berkeley, CA: North Atlantic Books.

Trauma Survivor's Inner Role Atom

Maté, G. (2008). *In the Realm of Hungry Ghosts: Close Encounters with Addiction*. Berkeley, CA: North Atlantic Books.

Moreno, J.L. (1946) *Psychodrama Volume 1*. Beacon, NY: Beacon House

Moreno, J.L. (1947). The Future of Man's World. *Psychodrama Monographs, No. 21*. Beacon, NY: Beacon House.

Moreno, J.L. (1953). *Who shall survive?* Beacon, NY: Beacon House.

Moreno, J.L. (1965). Therapeutic Vehicles and the Concept of Surplus Reality. *Group Psychotherapy* 18: 211-216

Moreno, Z.T. (2012). *To Dream Again: A Memoir*. New York: Mental Health Resources.

Moreno, Z. T., Blomkvist, L.D., & Rützel, E. (2000). *Psychodrama, surplus reality and the art of healing*. London, UK: Routledge Press.

Ogden, P., Minton, K., & Pain, C. (2006). *Trauma and the Body: A Sensorimotor Approach to Psychotherapy*. Norton Series on Interpersonal Neurobiology. New York, NY: WW Norton & Company.

Panksepp, J. and Biven, L. (2012). *The Archaeology of Mind: Neuroevolutionary Origins of Human Emotions*. New York, NY: W.W. Norton & Company, Inc.

Payne, P., Levine, P.A., & Crane-Godreau, M.A. (2015). Somatic experiencing: using interoception and proprioception as core elements of trauma therapy. *Frontiers in Psychology* 6: 93

Trauma Survivor's Inner Role Atom

Perry, R., Saby, K., Wenos, J., Hudgins, K., & Baller, S. (2016) Psychodrama Intervention for Female Service Members Using the Therapeutic Spiral Model. *The Journal of Psychodrama, Sociometry, and Group Psychotherapy*, 64(1): 11-23

Porges, S. W. (2017). *The Pocket Guide to The Polyvagal Theory: The Transformative Power of Feeling Safe*. New York, NY: W.W. Norton & Company

Schore, A. N. (2012). *The science of the art of psychotherapy*. New York, NY: W. W. Norton & Company.

Schmalzl, L., Powers, C., & Blom, E.H. (2015). Neurophysiological and neurocognitive mechanisms underlying the effects of yoga-based practices: towards a comprehensive theoretical framework. *Frontiers in Human Neuroscience* 9: 235

Siegel, D. J. (2006). An interpersonal neurobiology approach to psychotherapy: Awareness, mirror neurons, and neural plasticity in the development of wellbeing. *Psychiatric Annals* 36(4), 247-258

Siegel, D. J. (2012). *The Developing Mind: How relationship and the brain interact to shape who we are (2nd Edition)*. New York, NY: Guilford Press.

Toscani, M. F., & Hudgins, M.K. (1995). *The trauma survivor's intrapsychic role atom*. Workshop Handout. Madison, WI: The Center for Experiential Learning.

Van der Kolk, B. (2014). *The body keeps the score: Brain, mind, and body in the healing of trauma*. New York, NY: Viking Press.

Wylie, M. S. (2004). The limits of talk: Bessel van der Kolk wants to transform the treatment of trauma. *Psychotherapy Networker*, (28) 30-41.